

ΠΑΡΑΡΤΗΜΑ 13

COVID-19 Information

Public health information (CDC)


Research information (NIH)

SARS-CoV-2 data (NCBI)

Prevention and treatment information (HHS)

Español

FULL TEXT LINKS



Randomized Controlled Trial Rev Mal Respir. 2018 Mar;35(3):264-268.
doi: 10.1016/j.rmr.2017.01.010. Epub 2018 Feb 1.

[Effect of a surgical mask on six minute walking distance]

[Article in French]

E Person ¹, C Lemerrier ¹, A Royer ¹, G Reyckler ²

Affiliations

PMID: 29395560 DOI: 10.1016/j.rmr.2017.01.010

Abstract

Introduction: Six minutes walking test (6MWT) is regularly used in pulmonology. To minimize the risk of cross-infection, some patients must wear surgical mask at rest and sometimes during exercise.

Aim of the study: To evaluate the effect of wearing a surgical mask during 6MWT in healthy subjects.

Material and method: It is a prospective study on 44 healthy subjects. After a first 6MWT for training, they performed randomly two 6MWT: with or without a surgical mask. Distance and dyspnea, heart rate and saturation variations were recorded.

Results: Distance was not modified by the mask ($P=0.99$). Dyspnea variation was significantly higher with surgical mask (+5.6 vs. +4.6; $P<0.001$) and the difference was clinically relevant. No difference was found for the variation of other parameters.

Conclusion: Wearing a surgical mask modifies significantly and clinically dyspnea without influencing walked distance.

Keywords: Dyspnoea; Dyspnée; Masque de soins; Surgical mask; Test de marche; Walking test.

Copyright © 2017 SPLF. Published by Elsevier Masson SAS. All rights reserved.

Related information

Cited in Books

MedGen

LinkOut – more resources

Full Text Sources

Elsevier Science

Masson (France)

Other Literature Sources

scite Smart Citations

Medical

ClinicalTrials.gov

MedlinePlus Health Information

Research Materials

NCI CPTC Antibody Characterization Program